Copyright

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract); that it is not under consideration for publication elsewhere; that its publication has been approved by all coauthors, if any, as well as by the responsible authorities at the institute where the work has been carried out; that, if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher and that the manuscript will not be published elsewhere in any language without the consent of the copyright holders.

All articles published in this journal are protected by copyright, which covers the exclusive rights to reproduce and distribute the article (e. g., as offprints), all translation rights as well as the rights to publish the article in any electronic form. No material published in this journal may be reproduced photographically or stored on microfilm, in electronic data bases, video disks, etc., without first obtaining written permission from the publisher.

The use of general descriptive names, trade names, trademarks, etc., in this publication, even if not specifically identified, does not imply that these names are not protected by the relevant laws and regulations.

While the advice and information in this journal is believed to be true and accurate at the date of its going to press, neither the authors, the editors, nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Special regulations for photocopies in the USA. Photocopies may be made for personal or in-house use beyond the limitations stipulated under Section 107 or 108 of U.S. Copyright Law, provided a fee is paid. All fees should be paid to the Copyright Clearance Center, Inc., 21 Congress Street, Salem, MA 01970, USA, stating the ISSN 0012-186X, the volume, and the first and last page numbers of each article copied. The copyright owner's consent does not include copying for general distribution, promotion, new works, or resale. In these cases, specific written permission must first be obtained from the publisher.

Springer-Verlag Berlin Heidelberg New York London Paris Tokyo Hong Kong Barcelona Budapest
Printed in Germany by aprinta Wemding
© Springer-Verlag Berlin Heidelberg 1999
Springer-Verlag GmbH & Co.KG, D-14197 Berlin, Germany

Contents

Full Instructions to authors

A 6

Editorials

Hopper J. L.: Is Type II (non-insulin-dependent) diabetes mellitus not so "genetic" after all? 125
Waldhäusl W.: The world of Diabetologia in 1999 1

Reviews

Ashcroft F. M., Gribble F. M.: ATP-sensitive K^+ channels and insulin secretion: their role in health and disease 903
Auwerx J.: PPARγ, the ultimate thrifty gene 1033
Bolli G. B., Di Marchi R. D., Park G. D., Pramming S., Koivisto V. A.: Insulin analogues and their potential in the management of diabetes mellitus 1151
Coleman R. A., Herrmann T. S.: Nutritional regulation of leptin in humans 639
Hopfner R. L., Gopalakrishnan V.: Endothelin: emerging role in diabetic vascular complications 1383
Leslie R. D. G., Atkinson M. A., Notkins A. L.: Autoantigens IA-2 and GAD in Type I (insulin-dependent) diabetes 13
Malaisse W. J.: Insulinotropic action of monosaccharide esters: therapeutic perspectives 286
McGarry J. D., Dobbins R. L.: Fatty acids, lipotoxicity and insulin secretion 128
Mogensen C. E.: Microalbuminuria, blood pressure and diabetic renal disease: origin and development of ideas 263
Schmetterer L., Wolzt M.: Ocular blood flow and associated functional deviations in diabetic retinopathy 387
Sima A. A. F., Sugimoto K.: Experimental diabetic neuropathy: an update 773
Tomlinson D. R.: Mitogen-activated protein kinases as glucose transducers for diabetic complications 1271
Zimmet P. Z.: Diabetes epidemiology as a tool to trigger diabetes research and care 499

Articles

Alm P., Ekström P., Henningsson R., Lundquist I.: Morphological evidence for the existence of nitric oxide and carbon monoxide pathways in the rat islets of Langerhans: An immunocytochemical and confocal microscopical study 978
Almind K., Frederiksen S. K., Bernal D., Hansen T., Ambye L., Urhammmer S., Ekstrøm C. T., Berglund L., Reneland R., Lithell H., White M. F., Van Obberghen E., Pedersen O.: Search for variants of the gene-promoter and the potential phosphotyrosine encoding sequence of the insulin receptor substrate-2 gene: evaluation of their relation with alterations in insulin secretion and insulin sensitivity 1244
Andreelli F., Laville M., Ducluzeau P.-H., Vega N., Vallier P., Khalfallah Y., Riou J.-P., Vidal H.: Defective regulation of phosphatidylinositol-3-kinase gene expression in skeletal muscle and adipose tissue of non-insulin-dependent diabetes mellitus patients 358
Balkan B., Kwansnik L., Miserendino R., Holst J. J., Li X.: Inhibition of dipeptidyl peptidase IV with NVP-DPP728 increases plasma GLP-1 (7–36 amide) concentrations and improves oral glucose tolerance in obese Zucker rats 1324
Barazzoni R., Zanetti M., Tiengo A., Tessari P.: Protein metabolism in glucagonoma (Short communication) 326

Copyright

Submission of a manuscript implies: that the work described has not been published before (except in the form of an abstract); that it is not under consideration for publication elsewhere; that its publication has been approved by all coauthors, if any, as well as by the responsible authorities at the institute where the work has been carried out; that, if and when the manuscript is accepted for publication, the authors agree to automatic transfer of the copyright to the publisher and that the manuscript will not be published elsewhere in any language without the consent of the copyright holders.

All articles published in this journal are protected by copyright, which covers the exclusive rights to reproduce and distribute the article (e. g., as offprints), all translation rights as well as the rights to publish the article in any electronic form. No material published in this journal may be reproduced photographically or stored on microfilm, in electronic data bases, video disks, etc., without first obtaining written permission from the publisher.

The use of general descriptive names, trade names, trademarks, etc., in this publication, even if not specifically identified, does not imply that these names are not protected by the relevant laws and regulations.

While the advice and information in this journal is believed to be true and accurate at the date of its going to press, neither the authors, the editors, nor the publisher can accept any legal responsibility for any errors or omissions that may be made. The publisher makes no warranty, express or implied, with respect to the material contained herein.

Special regulations for photocopies in the USA. Photocopies may be made for personal or in-house use beyond the limitations stipulated under Section 107 or 108 of U.S. Copyright Law, provided a fee is paid. All fees should be paid to the Copyright Clearance Center, Inc., 21 Congress Street, Salem, MA 01970, USA, stating the ISSN 0012-186X, the volume, and the first and last page numbers of each article copied. The copyright owner's consent does not include copying for general distribution, promotion, new works, or resale. In these cases, specific written permission must first be obtained from the publisher.

Springer-Verlag Berlin Heidelberg New York London Paris Tokyo Hong Kong Barcelona Budapest
Printed in Germany by aprinta Wemding
© Springer-Verlag Berlin Heidelberg 1999
Springer-Verlag GmbH & Co.KG, D-14197 Berlin, Germany

Contents

Full Instructions to authors

A 6

Editorials

Hopper J. L.: Is Type II (non-insulin-dependent) diabetes mellitus not so "genetic" after all? 125
Waldhäusl W.: The world of Diabetologia in 1999 1

Reviews

Ashcroft F. M., Gribble F. M.: ATP-sensitive K^+ channels and insulin secretion: their role in health and disease 903
Auwerx J.: PPARγ, the ultimate thrifty gene 1033
Bolli G. B., Di Marchi R. D., Park G. D., Pramming S., Koivisto V. A.: Insulin analogues and their potential in the management of diabetes mellitus 1151
Coleman R. A., Herrmann T. S.: Nutritional regulation of leptin in humans 639
Hopfner R. L., Gopalakrishnan V.: Endothelin: emerging role in diabetic vascular complications 1383
Leslie R. D. G., Atkinson M. A., Notkins A. L.: Autoantigens IA-2 and GAD in Type I (insulin-dependent) diabetes 13
Malaisse W. J.: Insulinotropic action of monosaccharide esters: therapeutic perspectives 286
McGarry J. D., Dobbins R. L.: Fatty acids, lipotoxicity and insulin secretion 128
Mogensen C. E.: Microalbuminuria, blood pressure and diabetic renal disease: origin and development of ideas 263
Schmetterer L., Wolzt M.: Ocular blood flow and associated functional deviations in diabetic retinopathy 387
Sima A. A. F., Sugimoto K.: Experimental diabetic neuropathy: an update 773
Tomlinson D. R.: Mitogen-activated protein kinases as glucose transducers for diabetic complications 1271
Zimmet P. Z.: Diabetes epidemiology as a tool to trigger diabetes research and care 499

Articles

Alm P., Ekström P., Henningsson R., Lundquist I.: Morphological evidence for the existence of nitric oxide and carbon monoxide pathways in the rat islets of Langerhans: An immunocytochemical and confocal microscopical study 978
Almind K., Frederiksen S. K., Bernal D., Hansen T., Ambye L., Urhammmer S., Ekstrøm C. T., Berglund L., Reneland R., Lithell H., White M. F., Van Obberghen E., Pedersen O.: Search for variants of the gene-promoter and the potential phosphotyrosine encoding sequence of the insulin receptor substrate-2 gene: evaluation of their relation with alterations in insulin secretion and insulin sensitivity 1244
Andreelli F., Laville M., Ducluzeau P.-H., Vega N., Vallier P., Khalfallah Y., Riou J.-P., Vidal H.: Defective regulation of phosphatidylinositol-3-kinase gene expression in skeletal muscle and adipose tissue of non-insulin-dependent diabetes mellitus patients 358
Balkan B., Kwansnik L., Miserendino R., Holst J. J., Li X.: Inhibition of dipeptidyl peptidase IV with NVP-DPP728 increases plasma GLP-1 (7–36 amide) concentrations and improves oral glucose tolerance in obese Zucker rats 1324
Barazzoni R., Zanetti M., Tiengo A., Tessari P.: Protein metabolism in glucagonoma (Short communication) 326

Böni-Schnetzler M., Hauri C., Zapf J.: Leptin is suppressed during infusion of recombinant human insulin-like growth factor I (rhIGF I) in normal rats 160
Boomsma F., Meiracker AH van den, Winkel S., Aanstoot H.J., Batstra M.R., Man in 't Veld A.J., Bruining G.J.: Circulating semicarbazide-sensitive amine oxidase is raised both in Type I (insulin-dependent), in Type II (non-insulin-dependent) diabetes mellitus and even in childhood Type I diabetes at first clinical diagnosis 233
Bosi E., Garancini M.P., Poggiali F., Bonifacio E., Gallus G.: Low prevalence of islet autoimmunity in adult diabetes and low predictive value of islet autoantibodies in the general adult population of northern Italy (Short communication) 840
Brock B., Gregersen S., Kristensen K., Thomsen J.L., Buschard K., Kofod H., Hermansen K.: The insulinotropic effect of endothelin-1 is mediated by glucagon release from the islet alpha cells 1302
Bruno G., Merletti F., Boffetta P., Cavallo-Perin P., Bargero G., Gallone G., Pagano G.: Impact of glycaemic control, hypertension and insulin treatment on general and cause-specific mortality: an Italian population-based cohort of Type II (non-insulin-dependent) diabetes mellitus (Short communication) 297
Buschard K., Schloot N.C., Kaas A., Bock T., Horn T., Fredman P., Roep B.O.: Inhibition of insulin-specific autoreactive T-cells by sulphatide which is variably expressed in beta cells 1212
Buteau J., Roduit R., Susini S., Prentki M.: Glucagon-like peptide-1 promotes DNA synthesis, activates phosphatidylinositol 3-kinase and increases transcription factor pancreatic and duodenal homeobox gene 1 (PDX-1) DNA binding activity in beta (INS-1)-cells (Rapid communication) 856
Cameron N.E., Cotter M.A., Jack A.M., Basso M.D., Hohman TC : Protein kinase C effects on nerve function, perfusion, Na^+, K^+-ATPase activity and glutathione content in diabetic rats 1120
Cassell P.G., Neverova M., Janmohamed S., Uwakwe N., Qureshi A., McCarthy M.I., Saker P.J., Albon L., Kopelman P., Noonan K., Easlick J., Ramachandran A., Snehalatha C., Pecqueur C., Ricquier D., Warden C., Hitman G.A.: An uncoupling protein 2 gene variant is associated with a raised body mass index but not Type II diabetes 688
Chang S.L., Lin J.G., Chi T.C., Liu I.M., Chen J.T.: An insulin-dependent hypoglycaemia induced by electroacupuncture at the Zhongwan (CV12) acupoint in diabetic rats 250
Chen M.-C., Schuit F., Eizirik D.L.: Indentification of IL-1β-induced messenger RNAs in rat pancreatic beta cells by differential display of messenger RNA 1199
Cherubini V., Carle F., Gesuita R., Iannilli A., Tuomilehto J., Prisco F., Iafusco D., Altobelli E., Chiarelli F., De Giorgi G., Falorni A.: Large incidence variation of Type I diabetes in central-southern Italy 1990–1995: lower risk in rural areas (Short communication) 789
Chowdhury T.A., Dyer P.H., Mijovic C.H., Dunger D.B., Barnett A.H., Bain S.C.: Human leucocyte antigen and insulin gene regions and nephropathy in Type I diabetes (Short communication) 1017
Dalgaard L.T., Sørensen T.I.A., Andersen T., Hansen T., Pedersen O.: An untranslated insertion variant in the uncoupling protein 2 gene is not related to body mass index and changes in body weight during a 26-year follow-up in Danish Caucasian men 1413
Davis E.A., Cuesta-Muñoz A., Raoul M., Buettger C., Sweet I., Moates M., Magnuson M.A., Matschinsky F.M.: Mutants of glucokinase cause hypoglycaemia- and hyperglycaemia syndromes and their analysis illuminates fundamental quantitative concepts of glucose homeostasis 1175
De Mattia G., Bravi M.C., Costanzo A., Laurenti O., Cassone Faldetta M., Armiento A., De Luca O., Ferri C.: Effects of insulin on in vitro vascular cell adhesion molecule-1 expression and in vivo soluble VCAM-1 release (Short communication) 1235
de Vegt F., Dekker J.M., Ruhé H.G., Stehouwer C.D.A., Nijpels G., Bouter L.M., Heine R.J.: Hyperglycaemia is associated with all-cause and cardiovascular mortality in the Hoorn population: the Hoorn Study 926
Deinum J., Rønn B., Mathiesen E., Derkx F.H.M., Hop W.C.J., Schalekamp MA.D.H.: Increase in serum prorenin precedes onset of microalbuminuria in patients with insulin-dependent diabetes mellitus 1006
Deng D., Evans T., Mukherjee K., Downey D., Chakrabarti S.: Diabetes-induced vascular dysfunction in the retina: role of endothelins 1228
Di Luca M., Ruts L., Gardoni F., Cattabeni F., Biessels G.J., Gispen W.H.: NMDA receptor subunits are modified transcriptionally and post-translationally in the brain of streptozotocin-diabetic rats 693
Douglas Petersen L., van der Keur M., de Vries R.R.P., Roep B.O.: Autoreactive and immunoregulatory T-cell subsets in insulin-dependent diabetes mellitus 443
Dupont S., Vionnet N., Chèvre J.C., Gallina S., Dina C., Seino Y., Yamada Y., Froguel P.: No evidence of linkage or diabetes-associated mutations in the transcription factors BETA2/NEUROD1 and PAX4 in Type II diabetes in France (Short communication) 480
Ebeling P., Teppo A.-M., Koistinen H.A., Viikari J., Rönnemaa T., Nissén M., Bergkulla S., Salmela P., Saltevo J., Veikko, Koivisto A.: Troglitazone reduces hyperglycaemia and selectively acute-phase serum proteins in patients with Type II diabetes 1433
Ek J., Urhammer S.A., Sørensen T.I.A., Andersen T., Auwerx J., Pedersen O.: Homozygosity of the *Pro12Ala* variant of the peroxisome proliferation-activated receptor-γ2 (*PPAR*-γ2): divergent modulating effects on body mass index in obese and lean Caucasian men (Short communication) 892
Elliott R.B., Harris D.P., Hill J.P., Bibby N.J., Wasmuth H.E.: Type I (insulin dependent) diabetes mellitus and cow milk: casein variant consumption 292
Eriksson J., Lindström J., Valle T., Aunola S., Hämäläinen H., Ilanne-Parikka P., Keinänen-Kiukaanniemi S., Laakso M., Lauhkonen M., Lehto P., Lehtonen A., Louheranta A., Mannelin M., Martikkala V., Rastas M., Sundvall J., Turpeinen A., Viljanen T., Uusitupa M., Tuomilehto J on behalf of the Finnish Diabetes Prevention Study Group: Prevention of Type II diabetes in subjects with impaired glucose tolerance: the Diabetes Prevention Study (DPS) in Finland. Study design and 1-year interim report on the feasibility of the lifestyle intervention programme 793
Esterbauer H., Oberkofler H., Dallinger G., Breban D., Hell E., Krempler F., Patsch W.: Uncoupling protein-3 gene expression: reduced skeletal muscle mRNA in obese humans during pronounced weight loss 302
Fagerudd J.A., Pettersson-Fernholm K.J., Grönhagen-Riska C., Groop P.-H.: The impact of a family history of Type II (non-insulin-dependent) diabetes mellitus on the risk of diabetic nephropathy in patients with Type I (insulin-dependent) diabetes mellitus 519
Fernández-Real J.-M., Ricart W.: Insulin resistance and inflammation in an evolutionary perspective: the contribution of cytokine genotype/phenotype to thriftiness 1367
Flück C.E., Kuhlmann B.V., Mullis P.E.: Insulin increases serum leptin concentrations in children and adolescents with newly diagnosed Type I diabetes mellitus with and without ketoacidosis (Short communication) 1067

Forouhi N.G., Jenkinson G., Thomas E.L., Mullick S., Mierisova S., Bhonsle U., McKeigue P.M., Bell J.D.: Relation of triglyceride stores in skeletal muscle cells to central obesity and insulin sensitivity in European and South Asian men (Short communication) 932

Forsblom C.M., Kanninen T., Lehtovirta M., Saloranta C., Groop L.C.: Heritability of albumin excretion rate in families of patients with Type II diabetes 1359

Forsén L., Meyer H.E., Midthjell K., Edna T.-H.: Diabetes mellitus and the incidence of hip fracture: results from the Nord-Trøndelag Health Survey 920

Furukawa M., Zhang Y.O., Nie L., Shibata H., Kojima I.: Role of mitogen-activated protein kinase and phosphoinositide 3-kinase in the differentiation of rat pancreatic AR42J cells induced by hepatocyte growth factor 450

Garofano A., Czernichow P., Bréant B.: Effect of ageing on beta-cell mass and function in rats malnourished during the perinatal period 711

Gautier J.-F., Milner M.R., Elam E., Chen K., Ravussin E., Pratley R.E.: Visceral adipose tissue is not increased in Pima Indians compared with equally obese Caucasians and is not related to insulin action or secretion 28

Ghosh S., Langefeld C.D., Ally D., Watanabe R.M., Hauser E.R., Magnuson V.L., Nylund S.J., Valle T., Eriksson J., Bergman R.N., Tuomilehto J., Collins F.S., Boehnke M.: The W64R variant of the β_3-adrenergic receptor is not associated with Type II diabetes or obesity in a large Finnish sample 238

Giannattasio C., Failla M., Piperno A., Grappiolo A., Gamba P., Paleari F., Mancia G.: Early impairment of large artery structure and function in Type I diabetes mellitus 987

Giltay E.J., Lambert J., Elbers J.M.H., Gooren L.J.G., Asscheman H., Stehouwer C.D.A.: Arterial compliance and distensibility are modulated by body composition in both men and women but by insulin sensitivity only in women 214

Giroix M.-H., Saulnier C., Portha B.: Decreased pancreatic islet response to L-leucine in the spontaneously diabetic GK rat: enzymatic, metabolic and secretory data 965

Gleeson A., Anderton K., Owens D., Bennett A., Collins P., Johnson A., White D., Tomkin G.H.: The Role of microsomal triglyceride transfer protein and dietary cholesterol in chylomicron production in diabetes 944

Goalstone M.L., Wall K., Leitner J.W., Kurowski T., Ruderman N., Pan S.J., Ivy J.L., Moller D.E., Draznin B.: Increased amounts of farnesylated p21Ras in tissues of hyperinsulinaemic animals 310

Greig N.H., Holloway H.W., De Ore K.A., Jani D., Wang Y., Zhou J., Garant M.J., Egan J.M.: Once daily injection of exendin-4 to diabetic mice achieves long-term beneficial effects on blood glucose concentrations 45

Gribble F.M., Ashcroft F.M.: Differential sensitivity of beta-cell and extrapancreatic K_{ATP} channels to gliclazide (Rapid communication) 845

Grill V., Persson G., Carlsson S., Norman A., Alvarsson M., Östensson C.-G., Svanström L., Efendic S.: Family history of diabetes in middle-aged Swedish men is a gender unrelated factor which associates with insulinopenia in newly diagnosed diabetic subjects 15

Hamilton-Wessler M., Ader M., Dea M., Moore D., Jorgensen P.N., Markussen J., Bergmann R.N.: Mechanism of protracted metabolic effects of fatty acid acylated insulin, NN304, in dogs: retention of NN304 by albumin 1254

Hammes H.-P., Alt A., Niwa T., Clausen J.T., Bretzel R.G., Brownlee M., Schleicher E.D.: Differential accumulation of advanced glycation end products in the course of diabetic retinopathy 728

Hammes H.-P., Brownlee M., Lin J., Schleicher E., Bretzel R.G.: Diabetic retinopathy risk correlates with intracellular concentrations of the glycoxidation product N^{ε}-(carboxymethyl) lysine independently of glycohaemoglobin concentrations (Short communication) 603

't Hart L.M., de Knijff P., Dekker J.M., Stolk R.P., Nijpels G., van der Does F.E.E., Ruige J.B., Grobbee D.E., Heine R.J., Maassen J.A.: Variants in the sulphonylurea receptor gene: association of the exon 16-3 *t* variant with Type II diabetes mellitus in Dutch Caucasians (Short communication) 617

Hermans M.P., Levy J.C., Morris R.J., Turner R.C.: Comparison of insulin sensitivity tests across a range of glucose tolerance from normal to diabetes 678

Hinokio Y., Suzuki S., Hirai M., Chiba M., Hirai A., Toyota T.: Oxidative DNA damage in diabetes mellitus: its association with diabetic complications (Short communication) 995

Hjeltnes N., Fernström M., Zierath J.R., Krook A.: Regulation of UCP2 and UCP3 by muscle disuse and physical activity in tetraplegic subjects 826

Hofmann M.A., Schiekofer S., Isermann B., Kanitz M., Henkels M., Joswig M., Treusch A., Morcos M., Weiss T., Borcea V., Abdel Khalek A.K.M., Amiral J., Tritschler H., Ritz E., Wahl P., Ziegler R., Bierhaus A., Nawroth P.P.: Peripheral blood mononuclear cells isolated from patients with diabetic nephropathy show increased activation of the oxidative-stress sensitive transcription factor NF-κB 222

Holemans K., Gerber R.T., Meurrens K., De Clerck F., Poston L., Van Assche F.A.: Streptozotocin diabetes in the pregnant rat induces cardiovascular dysfunction in adult offspring 81

Holness M.J., Sugden M.C.: In late pregnancy insulin-dependent glucose transport/phosphorylation is selectively impaired and activation of glycogen synthase by insulin facilitated in skeletal muscles of 24-h starved rats 802

Holzäpfel A., Festa A., Stacher-Janotta G., Bergmann H., Shnawa N., Brannath W., Schernthaner G., Stacher G.: Gastric emptying in non-insulin-dependent diabetes before and after therapy readjustment: no influence of actual blood-glucose concentration 1383

Hoorens A., Pipeleers D.: Nicotinamide protects human beta cells against chemically-induced necrosis, but not against cytokine-induced apoptosis 55

Höppener J.W.M., Oosterwijk C., Nieuwenhuis M.G., Posthuma G., Thijssen J.H.H., Vroom ThM, Ahrén B., Lips C.J.M.: Extensive islet amyloid formation is induced by development of Type II diabetes mellitus and contributes to its progression: pathogenesis of diabetes in a mouse model 427

Horton V., Stratton I., Bottazzo G.F., Shattock M., Mackay I., Zimmet P., Manley S., Holman R., Turner R., for the UK Prospective Diabetes Study (UKPDS) Group: Genetic heterogeneity of autoimmune diabetes: age of presentation in adults is influenced by HLA *DRB1* and *DQB1* genotypes (UKPDS 43) 608

Hou X., Ling Z., Quartier E., Foriers A., Schuit F., Pipeleers D., Van Schravendijk C.: Prolonged exposure of pancreatic beta cells to raised glucose concentrations results in increased cellular content of islet amyloid polypeptide precursors 188

Ibrahim S., Harris N.D., Radatz M., Selmi F., Rajbhandari S., Brady L., Jakubowski J., Ward J.D.: A new minimally invasive technique to show nerve ischaemia in diabetic neuropathy 737

Imagawa A., Hanafusa T., Itoh N., Waguri M., Yamamoto K., Miyagawa J., Moriwaki M., Yamagata K., Iwahashi H., Sada M., Tsuji T., Tamura S., Kawata S., Kuwajima M., Nakajima H., Namba M., Matsuzawa Y.: Immunological abnormalities in islets at diagnosis paralleled further deterioration of glycaemic control in patients with recent-onset Type I (insulin-dependent)

diabetes mellitus (Short communication) 574
Imanishi M., Yoshioka K., Konishi Y., Okumura M., Okada N., Sato T., Tanaka S., Fujii S., Kimura G.: Glomerular hypertension as one cause of albuminuria in Type II diabetic patients 999
Inoue S., Fukuda K., Kudara T., Igura T., Inui Y., Tamura S., Hanafusa T., Kawata S., Matsuzawa Y.: Augmented growth response to IGF-1 via increased IRS-1 in Chinese hamster ovary cells expressing kinase-negative insulin receptors 763
Ishiyama-Shigemoto S., Yamada K., Yuan X., Ichikawa F., Nonaka K.: Association of polymorphisms in the β2-adrenergic receptor gene with obesity, hypertriglyceridaemia, and diabetes mellitus (Short Communication) 98
Jacob S., Hauer B., Becker R., Artzner S., Grauer P., Löblein K., Nielsen M., Renn W., Rett K., Wahl H.-G., Stumvoll M., Häring H.-U.: Lipolysis in skeletal muscle is rapidly regulated by low physiological doses of insulin (Short communicaton) 1171
Jones S.C., Saunders H.J., Qi W., Pollock C.A.: Intermittent high glucose enhances cell growth and collagen synthesis in cultured human tubulointerstitial cells 1113
Jönsson B., Cook J.R., Pedersen T.R.: The cost-effectiveness of lipid lowering in patients with diabetes: results from the 4S trial 1293
Josefsen K., Sørensen L.R., Buschard K., Birkenbach M.: Glucose induces early growth response gene (Egr-1) expression in pancreatic beta cells 195
Jude E.B., Boulton A.J.M., Ferguson M.W.J., Appleton I.: The role of nitric oxide synthase isoforms and arginase in the pathogenesis of diabetic foot ulcers: possible modulatory effects by transforming growth factor beta 1 748
Kajimoto Y., Matsuoka T., Kaneto H., Watada H., Fujitani Y., Kishimoto M., Sakamoto K., Matsuhisa M., Kawamori R., Yamasaki Y., Hori M.: Induction of glycation suppresses glucokinase gene expression in HIT-T15 cells 1417
Kallmann B.A., Lampeter E.F., Hanifi-Moghaddam P., Hawa M., Leslie R.D.G., Kolb H.: Cytokine secretion patterns in twins discordant for Type I diabetes (Rapid communication) 1080
Kaneto H., Kajimoto Y., Fujitani Y., Matsuoka T., Sakamoto K., Matsuhisa M., Yamasaki Y., Hori M.: Oxidative stress induces p21 expression in pancreatic islet cells: possible implication in beta-cell dysfunction (Rapid communication) 1093
Kasuya Y., Ito M., Nakamura J., Hamada Y., Nakayama M., Chaya S., Komori T., Naruse K., Nakashima E., Kato K., Koh N., Hotta N.: An aldose redutase inhibitor prevents the intimal thickening in coronary arteries of galactose-fed beagle dogs 1404
Kawano Y., Rincon J., Soler A., Ryder J.W., Nolte L.A., Zierath J.R., Wallberg-Heriksson H.: Changes in glucose transport and protein kinase $C\beta_2$ in rat skeletal muscle induced by hyperglycaemia 1071
Keegan A., Cotter M.A., Cameron N.E.: Effects of diabetes and treatment with the antioxidant α-lipoic acid on endothelial and neurogenic responses of corpus cavernosum in rats 343
Kim A., Jun H.-S., Wong L., Stephure D., Pacaud D., Trussell R.A., Yoon J.-W.: Human endogenous retrovirus with a high genomic sequence homology with *IDDMK$_{1,2}$22* is not specific for Type I (insulin-dependent) diabetic patients but ubiquitous (Rapid communication) 413
Klein H.H., Müller R., Vestergaard H., Pedersen O.: Implications of compound heterozygous insulin receptor mutations in congenital muscle fibre type disproportion myopathy for the receptor kinase activation (Short communication) 245
Knott R.M., Robertson M., Muckersie E., Folefac V.A., Fairhurst F.E., Wileman S.M., Forrester J.V.: A model system for the study of human retinal angiogenesis: activation of monocytes and endothelial cells and the association with the expression of the monocarboxylate transporter type 1 (MCT-1) 870
Koch M., Rett K., Maerker E., Volk A., Haist K., Deninger M., Renn W., Häring H.U.: The PPARγ2 amino acid polymorphism Pro 12 Ala is prevalent in offspring of Type II diabetic patients and is associated to increased insulin sensitivity in a subgroup of obese subjects (Rapid communication) 758
Kohner E.M., Stratton I.M., Aldington S.J., Turner R.C., Matthews D.R.: Microaneurysms in the development of diabetic retinopathy (UKPDS 42) 1107
Kong M.-F., King P., Macdonald I.A., Blackshaw P.E., Horowitz M., Perkins A.C., Armstrong E., Buchanan K.D., Tattersall R.B.: Euglycaemic hyperinsulinaemia does not affect gastric emptying in Type I and Type II diabetes mellitus 365
Kronenberg F., Auinger M., Trenkwalder E., Irsigler K., Utermann G., Dieplinger H.: Is apolipoprotein(a) a susceptibility gene for Type I diabetes mellitus and related to long-term survival? 1021
Kunt T., Schneider S., Pfützner A., Goitum K., Engelbach M., Schauf B., Beyer J., Forst T.: The effect of human proinsulin C-peptide on erythrocyte deformability in patients with Type I diabetes mellitus 465
Lalain S., Chaillous L., Gouin E., Saï P.: Intensity and mechanisms of in vitro xenorecognition of adult pig pancreatic islet cells by CD_4^+ and CD_8^+ lymphocytes from Type I diabetic or healthy subjects (Short communication) 330
Lam K.S.L., Ma O.C.K., Wat N.M.S., Chan L.C., Janus E.D.: β-fibrinogen gene G/A-455 polymorphism in relation to fibrinogen concentrations and ischaemic heart disease in Chinese patients with Type II diabetes (Short communication) 1250
Lampasona V., Bonfanti R., Bazzigaluppi E., Venerando A., Chiumello G., Bosi E., Bonifacio E.: Antibodies to tissue transglutaminase C in Type I diabetes (Short communication) 1195
Larsen J.J.S., Dela F., Madsbad S., Galbo H.: The effect of intense exercise on postprandial glucose homeostasis in Type II diabetic patients 1282
Larsson H., Ahrén B.: Insulin resistant subjects lack islet adaptation to short-term dexamethasone-induced reduction in insulin sensitivity 936
Lefrandt J.D., Hoogenberg K., van Roon A.M., Dullaart R.P.F., Gans R.O.B., Smit A.J.: Baroreflex sensitivity is depressed in microalbuminuric Type I diabetic patients at rest and during sympathetic manoeuvres 1345
Lehto M., Wipemo C., Ivarsson S.-A., Lindgren C., Lipsanen-Nyman M., Weng J., Wirbell L., Widén E., Tuomi T., Groop L.: High frequency of mutations in MODY and mitochondrial genes in Scandinavian patients with familial early-onset diabetes 1131
Lenhard J.M., Lancaster M.E., Paulik M.A., Weiel J.E., Binz J.G., Sundseth S.S., Gaskill B.A., Lightfoot R.M., Brown H.R.: The RXR agonist LG100268 causes hepatomegaly, improves glycaemic control and decreases cardiovascular risk and cachexia in diabetic mice suffering from pancreatic beta-cell dysfunction 545
Li L., Oshida Y., Kusunoki M., Yamanouchi K., Johansson B.-L., Wahren J., Sato Y: Rat C peptide I and II stimulate glucose utilization in STZ-induced diabetic rats 958
Lindberg B., Ivarsson S.-A., Landin-Olsson M., Sundkvist G., Svanberg L., Lernmark Å: Islet autoantibodies in cord blood from children who developed Type I (insulin-dependent) diabetes mellitus before 15 years of age 181
Lindblad P., Chow W.H., Chan J., Bergström A., Wolk A., Gridley G., McLaughlin J.K., Nyrén O., Adami

H.O.: The role of diabetes mellitus in the aetiology of renal cell cancer 107
Linderkamp O., Ruef P., Zilow E.P., Hoffmann G.F.: Impaired deformability of erythrocytes and neutrophils in children with newly diagnosed insulin-dependent diabetes mellitus 865
MacArthur D.L.A., de Koning E.J.P., Verbeek J.S., Morris J.F., Clark A.: Amyloid fibril formation is progressive and correlates with beta-cell secretion in transgenic mouse isolated islets 1219
Maegawa H., Obata T., Shibata T., Fujita T., Ugi S., Morino K., Nishio Y., Kojima H., Hidaka H., Haneda M., Yasuda H., Kikkawa R., Kashiwagi A.: A new antidiabetic agent (JTT-501) rapidly stimulates glucose disposal rates by enhancing insulin signal transduction in skeletal muscle 151
Maggi D., Barbetti F., Cordera R.: Role of proline 193 in the insulin receptor post-translational processing 435
Mäkimattila S., Nikkilä K., Yki-Järvinen H.: Causes of weight gain during insulin therapy with and without metformin in patients with Type II diabetes mellitus 406
Mathews C.E., McGraw R.A., Dean R., Berdanier C.D.: Inheritance of a mitochondrial DNA defect and impaired glucose tolerance in BHE/Cdb rats 35
McKenna K., Morris A.D., Azam H., Newton R.W., Baylis P.H., Thompson C.J.: Exaggerated vasopressin secretion and attenuated osmoregulated thirst in human survivors of hyperosmolar coma (Short communication) 534
Medici F., Hawa M., Ianari A., Pyke D.A., Leslie R.D.G.: Concordance rate for Type II diabetes mellitus in monozygotic twins: actuarial analysis 146
Meyer T.W., Bennett P.H., Nelson R.G.: Podocyte number predicts long-term urinary albumin excretion in Pima Indians with Type II diabetes and microalbuminuria (Short communication) 1341
Moczulski D.K., Burak W., Doria A., Zychma M., Zukowska-Szczechowska E., Warram J.H., Grzeszczak W.: The role of aldose reductase gene in the susceptibility to diabetic nephropathy in Type II (non-insulin-dependent) diabetes mellitus (Short Communication) 94
Møller A.M., Ek J., Durviaux S.M., Urhammer S.A., Clausen J.O., Eiberg H., Hansen T., Rousseau G.G., Lemaigre F.P., Pedersen O.: Hepatocyte nuclear factor-6: associations between genetic variability and Type II diabetes and between genetic variability and estimates of insulin secretion (Short communication) 896
Møller A.M., Ek J., Durviaux S.M., Urhammer S.A., Clausen J.O., Eiberg H., Hansen T., Rousseau G.G., Lemaigre F.P., Pedersen O.: Hepatocyte nuclear factor-6: associations beween genetic variability and Type II diabetes and between genetic variability and estimates of insulin secretion 1011
Moriwaki M., Itoh N., Miyagawa J., Yamamoto K., Imagawa A., Yamagata K., Iwahashi H., Nakajima H., Namba M., Nagata S., Hanafusa T., Matsuzawa Y.: Fas and Fas ligand expression in inflamed islets in pancreas sections of patients with recent-onset Type I diabetes mellitus 1332
Morrissette D.L., Goldstein M.K., Raskin D.B., Rowland D.L.: Finger and penile tactile sensitivity in sexually functional and dysfunctional diabetic men 336
Movassat J., Portha B.: Beta-cell growth in the neonatal Goto-Kakisaki rat and regeneration after treatment with streptozotocin at birth 1098
Mykkänen L., Zaccaro D.J., Hales C.N., Festa A., Haffner S.M.: The relation of proinsulin and insulin to insulin sensitivity and acute insulin response in subjects with newly diagnosed Type II diabetes: the Insulin Resistance Atherosclerosis Study 1060
Neele D.M., de Wit E.C.M., Princen H.M.G.: Insulin suppresses apolipoprotein(a) synthesis by primary cultures of cynomolgus monkey hepatocytes (Short Communication) 41
Nelson R.G., Tan M., Beck G.J., Bennett P.H., Knowler W.C., Mitch W.E., Blouch K., Myers B.D.: Changing glomerular filtration with progression from impaired glucose tolerance to Type II diabetes mellitus (Short Communication) 90
Nielsen S., Schmitz A., Bacher T., Rehling M., Ingerslev J., Mogensen C.E.: Transcapillary escape rate and albuminuria in Type II diabetes. Effects of short-term treatment with low-molecular weight heparin (Short Communication) 60
Nyholm B., Walker M., Gravholt C.H., Shearing P.A., Sturis J., Alberti K.G.M.M., Holst J.J., Schmitz O.: Twenty-four-hour insulin secretion rates, circulating concentrations of fuel substrates and gut incretin hormones in healthy offspring of Type II (non-insulin-dependent) diabetic parents: evidence of several aberrations 1314
Obrosova I.G., Fathallah L., Lang H.J., Greene D.A.: Evaluation of a sorbitol dehydrogenase inhibitor on diabetic peripheral nerve metabolism: a prevention study 1187
Okubo M., Watanabe H., Fujikawa R., Kawamura T., Egusa G., Yamakido M.: Reduced prevalence of diabetes according to 1997 American Diabetes Association criteria (Short communication) 1168
Onkamo P., Väänänen S., Karvonen M., Tuomilehto J.: Worldwide increase in incidence of Type I diabetes – the analysis of the data on published incidence trends 1395
Orho-Melander M., Almgren P., Kanninen T., Forsblom C., Groop L.C.: A paired-sibling analysis of the XbaI polymorphism in the muscle glycogen synthase gene 1138
Pieper G.M.: Enhanced, unaltered and impaired nitric oxide-mediated endothelium-dependent relaxation in experimental diabetes mellitus: importance of disease duration 204
Poulsen P., Kyvik K.O., Vaag A., Beck-Nielsen H.: Heritability of Type II (non-insulin-dependent) diabetes mellitus and abnormal glucose tolerance – a population-based twin study 139
Reed M.J., Meszaros K., Entes L.J., Claypool M.D., Pinkett J.G., Brignetti D., Luo J., Khandwala A., Reaven G.M.: Effect of masoprocol on carbohydrate and lipid metabolism in a rat model of Type II diabetes 102
Riaz S., Malcangio M., Miller M., Tomlinson D.R.: A vitamin D_3 derivative (CB1093) induces nerve growth factor and prevents neurotrophic deficits in streptozotocin-diabetic rats 1308
Rolandsson O., Hägg E., Hampe C., Sullivan Jr E.P., Nilsson M., Jansson G., Hallmans G., Lernmark Å: Glutamate decarboxylase (GAD65) and tyrosine phosphatase-like protein (IA-2) autoantibodies index in a regional population is related to glucose intolerance and body mass index (Short communication) 555
Rondinone C.M., Carvalho E., Wesslau C., Smith U.P.: Impaired glucose transport and protein kinase B activation by insulin, but not okadaic acid, in adipocytes from subjects with Type II diabetes mellitus (Rapid communication) 819
Rosenbauer J., Herzig P., von Kries R., Neu A., Giani G.: Temporal, seasonal, and geographical incidence patterns of Type I diabetes mellitus in children under 5 years of age in Germany (Short communication) 1055
Rudberg S., Østerby R., Bangstad H.-J., Dahlquist G., Persson B.: Effect of angiotensin converting enzyme inhibitor or beta blocker on glomerular structural changes in young microalbuminuric patients with Type I (insulin-dependent) diabetes mellitus 589
Rudich A., Tirosh A., Potashnik R., Khamaisi M., Bashan N.: Lipoic acid protects against oxidative stress induced impairment in insulin stimula-

tion of protein kinase B und glucose transport in 3T3-L1 adipocytes 949
Ruige J. B., Dekker J. M., Nijpels G., Popp-Snijders C., Stehouwer C. D. A., Kostense P. J., Bouter L. M., Heine R. J.: Hyperproinsulinaemia in impaired glucose tolerance is associated with a delayed insulin response to glucose (Short communication) 177
Salas A., Panés J., Rosenbloom C. L., Elizalde J. I., Anderson D. C., Granger D. N., Piqué J. M.: Differential effects of a nitric oxide donor on reperfusion-induced microvascular dysfunction in diabetic and non-diabetic rats 1350
Sánchez-Margalet V.: Modulation of insulin receptor signalling by pancreastatin in HTC hepatoma cells 317
Sánchez-Soto M. C., Larrieta M. E., Vidaltamayo R., Hiriart M.: Heterogeneity in glutamic acid decarboxylase expression among single rat pancreatic beta cells 1086
Sato A., Tarnow L., Parving H.-H.: Prevalence of left ventricular hypertrophy in Type I diabetic patients with diabetic nephropathy 76
Schaeffer G., Wascher T. C., Kostner G. M., Graier W. F.: Alterations in platelet Ca^{2+} signalling in diabetic patients is due to increased formation of superoxide anions and reduced nitric oxide production 167
Schalkwijk C. G., Poland D. C. W., Dijk W van, Kok A., Emeis J. J., Dräger A. M., Doni A., Hinsbergh VWM van, Stehouwer C. D. A.: Plasma concentration of C-reactive protein is increased in Type I diabetic patients without clinical macroangiopathy and correlates with markers of endothelial dysfunction: evidence for chronic inflammation 351
Schenker M., Hummel M., Ferber K., Walter M., Keller E., Albert E. D., Janka H.-U., Kastendiek C., Sorger M., Louwen F., Ziegler A.-G.: Early expression and high prevalence of islet autoantibodies for DR3/4 heterozygous and DR4/4 homozygous offspring of parents with Type I diabetes: The German BABYDIAB study 671
Schlingemann R. O., Hofman P., Vrensen GF. J. M., Blaauwgeers H. G. T.: Increased expression of endothelial antigen PAL-E in human diabetic retinopathy correlates with microvascular leakage 596
Shaw J. E., Hodge A. M., de Courten M., Chitson P., Zimmet P. Z.: Isolated post-challenge hyperglycaemia confirmed as a risk factor for mortality 1050
Shaw J. T. E., Purdie D. M., Neil H. A. W., Levy J. C., Turner R. C.: The relative risks of hyperglycaemia, obesity and dyslipidaemia in the relatives of patients with Type II diabetes mellitus (Short Communication) 24
Soulis T., Sastra S., Thallas V., Mortensen S. B., Wilken M., Clausen J. T., Bjerrum O. J., Petersen H., Lau J., Jerums G., Boel E., Cooper M. E.: A novel inhibitor of advanced glycation end-product formation inhibits mesenteric vascular hypertrophy in experimental diabetes 472
Strebelow M., Schlosser M., Ziegler B., Rjasanowski I., Ziegler M.: Karlsburg Type I diabetes risk study of a general population: frequencies and interactions of the four major Type I diabetes-associated autoantibodies studied in 9419 schoolchildren 661
Strobel A., Siquier K., Zilberfarb V., Strosberg A. D., Issad T.: Effect of thiazolidinediones on expression of UCP2 and adipocyte markers in human PAZ6 adipocytes 527
Sugimoto H., Shikata K., Wada J., Horiuchi S., Makino H.: Advanced glycation end products-cytokine-nitric oxide sequence pathway in the development of diabetic nephropathy: aminoguanidine ameliorates the overexpression of tumour necrosis factor-α and inducible nitric oxide synthase in diabetic rat glomeruli 878
Suschek C. V., Bonmann E., Kolb-Bachofen V.: A regulatory defect of constitutive no-synthase in islet endothelial cells correlates with probability of disease manifestation in BBdp rats 457
Suzuki M., Fujikura K., Kotake K., Inagaki N., Seino S., Takata K.: Immunolocalization of sulphonylurea receptor 1 in rat pancreas 1204
Tanaka T., Itoh H., Doi K., Fukunaga Y., Hosoda K., Shintani M., Yamashita J., Chun T.-H., Inoue M., Masatsugu K., Sawada N., Saito T., Inoue G., Nishimura H., Yoshimasa Y., Nakao K.: Down regulation of peroxisome proliferator-activated receptorγ expression by inflammatory cytokines and its reversal by thiazolidinediones 702
Tanizawa Y., Ohta Y., Nomiyama J., Matsuda K., Tanabe K., Inoue H., Matsutani A., Okuya S., Oka Y.: Overexpression of dominant negative mutant hepatocyte nuclear factor (HNF)-1α inhibits arginine-induced insulin secretion in MIN6 cells (Short communication) 887
Tannous M., Rabini R. A., Vignini A., Moretti N., Fumelli P., Zielinski B., Mazzanti L., Mutus B.: Evidence for *i*NOS-dependent peroxynitrite production in diabetic platelets 539
The DECODE-study group on behalf of the European Diabetes Epidemiology Group: Is fasting glucose sufficient to define diabetes? Epidemiological data from 20 European studies 647
The EURODIAB Substudy 2 Study Group: Vitamin D supplement in early childhood and risk for Type I (insulin-dependent) diabetes mellitus 51
Thorburn A. W., Baldwin M. E., Rosella G., Zajac J. D., Fabris S., Song S., Proietto J.: Features of syndrome X develop in transgenic rats expressing a non-insulin responsive phosphoenolpyruvate carboxykinase gene 419
Tiedge M., Lortz S., Munday R., Lenzen S.: Protection against the co-operative toxicity of nitric oxide and oxygen free radicals by overexpression of antioxidant enzymes in bioengineered insulin-producing RINm5F cells 849
Trovati M., Massucco P., Mattiello L., Costamagna C., Aldieri E., Cavalot F., Anfossi G., Bosia A., Ghigo D.: Human vascular smooth muscle cells express a constitutive nitric oxide synthase that insulin rapidly activates, thus increasing guanosine 3′ : 5′-cyclic monophosphate and adenosine 3′ : 5′-cyclic monophosphate concentrations 831
Tsuchida K., Makita Z., Yamagishi S., Atsumi T., Miyoshi H., Obara S., Ishida M., Ishikawa S., Yasumura K., Koike T.: Suppression of transforming growth factor beta and vascular endothelial growth factor in diabetic nephropathy in rats by a novel advanced glycation end product inhibitor, OPB-9195 579
Tuomilehto J., Karvonen M., Pitkäniemi J., Virtala E., Kohtamäki K., Toivanen L., Tuomilehto-Wolf E., The Finnish Childhood Type I Diabetes Registry Group: Record-high incidence of Type I (insulin-dependent) diabetes mellitus in Finnish children 655
van den Ouweland J. M. W., Maechler P., Wollheim C. B., Attardi G., Maassen J. A.: Functional and morphological abnormalities of mitochondria harbouring the $tRNA^{Leu(UUR)}$ mutation in mitochondrial DNA derived from patients with maternally inherited diabetes and deafness (MIDD) and progressive kidney disease 485
Varadi A., Lebel L., Hashim Y., Metha Z., Ashcroft S. J. H., Turner R.: Sequence variants of the sarco(endo)-plasmic reticulum Ca^{2+}-transport ATPase 3 gene (SERCA3) in Caucasian Type II diabetic patients (UK Prospective Diabetes Study 48) (Short communication) 1240
Veglio M., Borra M., Stevens L. K., Fuller J. H., Perin PC and the EURODIAB IDDM Complications Study Group: The relation between QTc interval prolongation and diabetic complications. The EURODIAB IDDM Complication Study Group 68
Wada R., Sugo M., Nakano M., Yagihashi S.: Only limited effects of aminoguanidine treatment on peripheral nerve function, (Na^+,K^+)-ATPase activity and thrombomodulin expression

in streptozotocin-induced diabetic rats 743
Wahren J., Ekberg K., Fernqvist-Forbes E., Nair S.: Brain substrate utilisation during acute hypoglycaemia 812
Wolf G., Wenzel U., Ziyadeh F. N., Stahl R. A. K.: Agiotensin converting-enzyme inhibitor treatment reduces glomerular $p16^{INK4}$ and $p27^{Kip1}$ expression in diabetic BBdp rats 1425
Xu X.-J., Gearon C., Stevens E., Vergani D., Baum H., Peakman M.: Spontaneous T-cell proliferation in the non-obese diabetic mouse to a peptide from the unique class II MHC molecule, I-A^{g7}, which is also protective against the development of autoimmune diabetes 560
Yamaoka T., Yano M., Idehara C., Yamada T., Tomonari S., Moritani M., Ii S., Yoshimoto K., Hata J., Itakura M.: Apoptosis and remodelling of beta cells by paracrine interferon-γ without insulitis in transgenic mice 566
Yoshiuchi I., Yamagata K., Yang Q., Iwahashi H., Okita K., Yamamoto K., Oue T., Imagawa A., Hamaguchi T., Yamasaki T., Horikawa Y., Satoh T., Nakajima H., Miyazaki J., Higashiyama S., Miyagawa J., Namba M., Hanafusa T., Matsuzawa Y.: Three new mutations in the hepatocyte nuclear factor-1 α gene in Japanese subjects with diabetes mellitus: clinical features and functional characterization (Short communication) 621
Zhang Y.-Q., Kanzaki M., Furukawa M., Shibata H., Ozeki M., Kojima I.: Involvement of Smad proteins in the differentiation of pancreatic AR42J cells induced by activin A 719

Rapid communication

Krssak M., Falk Petersen K., Dresner A., DiPietro L., Vogel S. M., Rothman D. L., Roden M., Shulman G. I.: Intramyocellular lipid concentrations are correlated with insulin sensitivity in humans: a ^{1}H NMR spectroscopy study 113

For debates

Gerbitz K.-D.: Reflexions on a newly discovered diabetogenic gene, wolframin (*WFS1*) 627
Nauck M. A.: Is glucagon-like peptide 1 an incretin hormone? 373

Letters

Arya S. C.: Temperature loggers are to be used to guarantee quality of insulin preparations 257
Asano T., Tsukuda K., Katagiri H., Onishi Y., Sakoda H., Ono H., Ogiliara T., Funaki M., Anai M., Inukai K., Fujishiro Y. M., Fukushima, Yamasoba T., Oka T., Kikuchi M., Oka Y.: Clinical relevance of heteroplasmic concentration of mitochondrial A 3243 G mutation in leucocytes 1439
Beales P. E., Pozzilli P.: Caution is required with Interferon α as a potential treatment for Type I diabetes 385
Bhonde R. R., Parab P. B.: Muscle insulin resistance: a probable case of disuse of skeletal muscles. Lamarckism revisited 257
Bloom S. R., Edwards C. M. B.: The incretins – outdated terminology in man? 1048
Boutin P., Gresh L., Cisse A., Hara M., Bell G., Babu S., Eisenbarth G., Froguel P.: Missense mutation Gly574Ser in the transcription factor HNF-1α is a marker of atypical diabetes mellitus in African-American children 380
Bring J., Wernroth L.: Inefficient analysis of twin data: is there an association between diabetes and birth weight? 898
Chico A., Pérez A., Córdoba A., Arcelús R., Carreras G., Leiva A de, González Sastre F., Blanco-Vaca F. 383
Cordain L., Miller J., Mann N.: Scant evidence of periodic starvation among hunter-gatherers 383
Cox R. D., Southam L., Hashim Y., Horton V., Mehta Z., Taghavi J., Lathrop M., Turner R.: UKPDS 31: Hepatocyte nuclear factor-1alpha (the MODY3 gene) mutations in late onset Type II diabetic patients in the United Kingdom 120
Cseh K., Baranyi É, Winkler G.: The role of cytokines of the innate and adaptive immune system in the regulation of insulin resistance 497
Dabelea D., Palmer J. P., Bennett P. H., Pettitt D. J., Knowler W. C.: Absence of glutamic acid decarboxylase antibodies in Pima Indian children with diabetes mellitus 1265
Dyer P. H., Chowdhury T. A., Dronsfield M. J., Dunger D., Barnett A. H., Bain S. C.: The 5'-end polymorphism of the aldose reductase gene is not associated with diabetic nephropathy in Caucasian Type I diabetic patients 1030
Faehling M., Fussgaenger R. D., Jehle P. M.: High affinity binding sites for proinsulin on human umbilical vein endothelial cells (HUVEC) 259
Grzeszczak W., Saucha W., Zychma M. J., Zukowska-Szczechowska E., Labuz B., Lacka B., Szydlowska I.: Is Trp64Arg polymorphism of β_3-adrenergic receptor a clinically useful marker for the predisposition to diabetic nephropathy in Type II diabetic patients? 632
Hanaire-Broutin H., Melki V., Tauber J. P.: Treatment of acquired immune insulin resistance with i. v. immunoglobulins 1378
Holmes-Walker D. J., Boyages S. C.: Prevalence of maternally inherited diabetes and deafness in Australian diabetic subjects 1028
Isotani H., Kameoka K., Nagano Y., Kitaoka H., Ohsawa N.: Characteristic neuroimaging findings in patients with diabetes and the 8296 mitochondrial $tRNA^{Lys}$ 1266
Karlsson S., Ahrén B.: Capsaicin-induced sensory denervation increases glucose elimination in rodents 260
Koehler C., Temelkova-Kurktschiev T., Henkel E., Schaper F., Fuecker K., Hanefeld M.: Is the newly suggested fasting plasma glucose cut-off point for the diagnosis of diabetes the right one? 635
Krssak M., Falk Petersen K., Dresner A., DiPietro L., Vogel S. M., Rothman D. L., Shulman G. I., Roden M.: Intramyocellular lipid concentrations are correlated with insulin sensitivity in humans: a ^{1}H NMR spectroscopy study (Rapid communication) 1269
Lauro D., Kido Y., Hayashi H., Ebina Y., Accili D.: Expression of kinase-inactive mutant insulin receptors does not rescue insulin receptor-deficient mice from perinatal death 1441
Lernmark Å, Kolb H., Mire-Sluis T.: Towards a World Health Organisation (WHO) approved standard sample for islet cell antibodies, GAD65 and IA-2 autoantibodies 381
Lindberg B., Ivarsson S.-A., Lernmark Å: Influence of heredity on the appearance of islet autoantibodies in cordblood from children developing Type I diabetes before 15 years of age 896
Lindberg B., Ivarsson S.-A., Lernmark Å: Islet autoantibodies in cord blood could be a risk factor for future diabetes 1375
Lindner T. H., Cockburn B. N., Bell G. I.: Molecular genetics of MODY in Germany 121
Malaisse W. J.: D-glucose transport and utilization in pituitary adenoma cells transfected with glucokinase gene 498
Mauricio D.: Islet autoantibodies at birth in children later developing Type I diabetes mellitus 896
McShane P.: Confidence intervals for frequencies of rare alleles 635
Naserke H. E., Schenker M., Ziegler A.-G.: Frequency of islet autoantibodies in relatives of patients with Type II diabetes or gestational diabetes 1377
Nauck M.: Response to: "The incretins – outdated terminology in man? By C. M. B. Edwards and S. R. Bloom, London UK" 1049

Nisoli E., Vettor R., Tonello C., Macor C., Federspil G., Carruba M. O.: Nutrient channelling-regulated peroxisome proliferator-activated receptor-γ-2 (PPARγ-2) and leptin gene expression in human subcutaneous fat 495
Odawara M., Yamashita K.: A common mutation of the methylenetetrahydrofolate reductase gene as a risk factor for diabetic nephropathy 631
Panarotto D., Maheux P.: Reduction of plasma leptin during a short-term fast, an oral glucose tolerance or a meal test can be a misleading bias in clinical studies 634
Pérez de Nanclares G., Bilbao J. R., Nisticò L., Buzzetti R., Larsen Z. M., Pociot F., Castaño L.: No evidence of association of chromosome 2q with Type I diabetes in the Basque population 119
Petrie J. R., Cleland S. J., Ueda S., Elliott H. L., Connell J. M. C.: Insulin-mediated vasodilation and forearm plethysmography: physiology not pharmacology 1376
Poulsen P., Vaag A. A., Kyvik K. O., Møller Jensen D., Beck-Nielsen H.: Evidence for an association between diabetes and birth weight still exists 899
Princen H. M. G.: Insulin suppresses apolipoprotein(a) synthesis in cultures cynomolgus monkey hepatocytes 897
Rathmann W., Giani G., Mielck A.: Cardiovascular risk factors in newly diagnosed abnormal glucose tolerance: comparison of 1997 ADA and 1985 WHO criteria 1268
Reaven G. M.: Insulin resistance, the key to survival: a rose by any other name 384
Riemens S. C., Dullaart R. P. F.: Effect of insulin on apolipoprotein(a) synthesis 897
Roep B. O., on behalf of the Immunology of Diabetes Society T-cell Committee: Standardization of T-cell assays in Type I diabetes 636
Seissler J., Schott M., Boms S., Wohlrab U., Ostendorf B., Morgenthater N. G., Scherbaum W. A.: Autoantibodies to human tissue transgiutaminase identify silent coeliac disease Type I diabetes 1440
Smith S. R., Lovejoy J. C., DeLaney J. P., Ruyter I., McNeil M.: The inhibition of glycogen synthesis by fatty acids is not mediated by inhibition of extracellular signal-regulated kinase 1/2 1031
Smulders Y. M., Brouwer C. B., Silberbusch J.: Is plasma homocysteine related to albumin excretion rate in patients with diabetes mellitus? 382
Steinberg H. O., Baron A. D.: Insulin-mediated vasodilation: why one's physiology could be the other's pharmacology 493
Stobbe I., Duinkerken G., van Rood J. J., Lernmark Å, Keymeulen B., Pipeleers D., De Vries R. R. P., Claas F. H. J., Roep B.: Tolerance to kidney allograf transplanted into Type I diabetic patients persists after in vivo challenge with pancreatic islet allografts that express repeated mismatches 1379
Wasada T., Katsumori K., Kuroki H., Iwamoto Y.: Insulin resistance facilitates the development of coronary artery disease in Japanese Type II diabetic patients: a single hospital-based follow-up study 1264
Weitgasser R., Davalli A. M., Weir G. C.: Measurement of glucose concentrations in rats: differences between glucose meter and plasma laboratory results 256
Worm D., Vinten J., Beck-Nielsen H.: The significance of phosphotyrosine phosphatase (PTPase) 1B in insulin signalling 1046
Yki-Järvinen H., Utriainen T.: Insulin-mediated vasodilation: the authors' reply 494
Young A., Kolterman O., Hall J.: Amylin innocent in essential hypertension? 1029
Zerbini G., Meregalli G., Cornaghi L., Mangili R., Pozza G.: Tetraploid cells and diabetic nephropathy 118
Zozuliñska D., Majchrzak A., Sobieska M., Wiktorowicz K., Wierusz-Wysocka B.: Serum interleukin-8 level is increased in diabetic patients 117

Abstracts

A 11

Organisation Section

A 355

Book review

901

EASD News Section

15, 21, 23, 27, 29, 33, 41, 51

Erratum

262, 386, 498, 1032, 1269, 1444

Author index

1453

Subject index

1458

Contents of Volume 42

I–IX